

Preliminary. Please do not cite/distribute without permission.

## **Mortality Among Persons Who Self-Identify as American Indian or Alaska Native**

Michael Bird (Kewa Pueblo)

Jacob Bor\* (Boston University)

Jahn Hakes (U.S. Census Bureau)

David Himmelstein (City University of New York)

Steffie Woolhandler (City University of New York)

[\\*jbor@bu.edu](mailto:*jbor@bu.edu)

*Prepared for NBER Racial and Ethnic Health Disparities Workshop, April 28, 2023*

### **ABSTRACT**

Race is not always reported accurately on death certificates, which can lead to biased estimates of mortality rates when numerators and denominators are taken from different sources. Native American/American Indian/Alaskan Native (AIAN) identity in particular is significantly underreported, leading to underestimates of mortality rates in this population. Using nationally representative, longitudinal data from the Mortality Disparities in American Communities study, 2008-2015, we estimate mortality rates and life expectancy for self-identified AIAN and other racial/ethnic groups without bias due to misclassification on death certificates. We find that the excess mortality risk among AIAN relative to the total U.S. population increases by 53% after accounting for misclassification.

## 1. Background

Life expectancy in the U.S. varies by race/ethnicity. Official U.S. government estimates rely on matching death certificates with population denominators. However, demographics on death certificates are not always reported accurately, which can lead to biased mortality estimates.<sup>1 2</sup> Native American/American Indian/Alaskan Native (AIAN) identity in particular is significantly underreported in death certificates,<sup>3</sup> despite rapid growth in the U.S. population identifying as AIAN on Census forms in recent decades.<sup>4</sup> The invisibility of AIAN in Census and health data has been referred to as “statistical erasure”<sup>5</sup> and follows centuries of deliberate policy to erase indigenous culture and identity, e.g. through genocide and forced assimilation.<sup>6</sup> The under-reporting of mortality among AIAN is particularly troubling because, even with the under-reporting, AIAN have among the highest mortality rates of any U.S. racial/ethnic group.

In this paper, we analyze restricted-use data from the Mortality Disparities in American Communities (MDAC) study. MDAC is a collaboration between the U.S. Census and National Vital Statistics System (NVSS). Specifically, the U.S. Census has linked individual respondents in the 2008 American Community Survey (ACS) to U.S. death records through 2015. (MDAC is in the process of being updated to 2019.) The MDAC database offers the opportunity to estimate population mortality rates based on 7 years of prospective follow-up for a representative population of the U.S. with different sociodemographic characteristics. In contrast to the conventional race-stratified mortality rates reported by CDC, the longitudinal mortality estimates from the MDAC are based on self-identified race/ethnicity, are not biased due to misreporting on death certificates.

---

<sup>1</sup> Arias E. The Validity of race and Hispanic origin reporting on death certificates in the United States. CDC 2008. Series: DHHS publication no. (PHS) 2009-1348. <https://stacks.cdc.gov/view/cdc/7066>

<sup>2</sup> Arias E, Heron MP, Hakes JK. Validity of race and Hispanic origin reporting on death certificates in the United States: an update. CDC. Vital and health statistics. Series 2, Data evaluation and methods research; no. 172, 2016. <https://stacks.cdc.gov/view/cdc/45533>

<sup>3</sup> *Op Cit.* Arias et al. 2014

<sup>4</sup> Passel JF. The growing American Indian Population: beyond demography. In: National Research Council, Committee on Population. Changing numbers, changing needs: American Indian demography and public health. 1996. <https://www.nap.edu/catalog/5355/changing-numbers-changing-needs-american-indian-demography-and-public-health>

<sup>5</sup> Small-Rodriguez D. <https://www.drdrdesi.com/research>

<sup>6</sup> Dunbar-Ortiz R. 2014. *An Indigenous People's History of the United States*.

Prior research has found substantial underreporting of AIAN racial identify on death certificates, ranging from 10% to 40%. Using death certificates augmented by Indian Health Service patient registration records, Arias et al<sup>7</sup> assessed AIAN life expectancy in 2007-2009 among persons living on or near tribal lands. They estimated AIAN life expectancy in the areas they studied at 71.1 years, compared to 78.4 years for the non-Hispanic White population nationally. A study of the AIAN population in the Pacific Northwest yielded similar findings.<sup>8</sup> Analyzing data on a sample of deaths linked to the 2010 Census, Arias and colleagues found that AIAN deaths are underreported by 25% in death certificates.<sup>9</sup> However, no recent longitudinal follow-up study has yet assessed AIAN mortality rates prospectively. The MDAC data enable calculation of unbiased estimates of AIAN age-specific mortality rates and life expectancy; comparison of mortality rates for different groups, e.g. AIAN alone, AIAN in combination with another race, Hispanic AIAN, AIAN on vs. off reservations; and assessment of individual-level socioeconomic factors as mediators of mortality disparities.

## 2. Data Description

### *Study population*

The study population consisted of all persons residing in the U.S. in 2008, the sampling frame for the 2008 ACS. The ACS is a national survey conducted by the U.S. Census, with data collected every month of the year in all states, D.C., and Puerto Rico. Each year the survey is sent to a 2.5% sample of the U.S. resident population. Surveys are filled out by mail or online. The Census Bureau follows up with a sample of non-responses by telephone and through in-person visits. The ACS includes both people residing in households and those residing in group quarters, such as schools, prisons, and nursing homes.

---

<sup>7</sup> Arias E, Xu J, Jim MA. Period life tables for the non-Hispanic American Indian and Alaska Native population, 2007–2009. *Am J Public Health*. 2014;104(S3):S312-9.

<sup>8</sup> Dankovchik J, Hoopes MJ, Warren-Mears V, Knaster E. Disparities in life expectancy of Pacific Northwest American Indians and Alaska natives: analysis of linkage-corrected life tables. *Public Health Reports*. 2015 Jan;130(1):71-80.

<sup>9</sup> Arias E, Xu J, Curtin S, Bastian B, Tejada-Vera B. 2021. Mortality Profile of the Non-Hispanic American Indian or Alaska Native Population, 2019. NVS Report 70(12).

Preliminary. Please do not cite/distribute without permission.

### *Data linkage*

The U.S. Census Bureau has linked the 2008 ACS at the individual level with mortality records from the National Death Index to construct the MDAC database. The MDAC was established to enable research on mortality disparities by sociodemographic and small-area geographic variables.<sup>10</sup> The MDAC includes information on date of death, place of death, and cause of death (ICD-10 codes). Linkage between the ACS and National Death Index was conducted using an exact match on Social Security Number (SSN) or Name and Date of Birth. 91% of records in the ACS have SSN, enabling highly accurate matches. The remainder are linked on name and birth date.<sup>11</sup> The linkage algorithm is tuned to be conservative such that a match would only be made on name and birth date if there was very high confidence it was the same person.

### *Access to the MDAC*

Access to the MDAC is restricted and we obtained approval to conduct this project from the Census. The team included a Census Statistician in the Census's Mortality Research Group who conducted the analyses and had direct access to the data. The broader team designed the analyses and Dr. Bor, who obtained Special Sworn Status, reviewed intermediate output through the Census's Research Output Direct Access System (RODAS). All output reported in this paper have been reviewed by the Census's Data Review Board (DRB) to ensure compliance with policies to protect confidentiality of respondents. Exact counts have been rounded according to a specific algorithm and small cells suppressed.

### *How representative is the MDAC?*

The MDAC is intended to include all respondents in the 2008 ACS. However, not all respondents had adequate identifiers (Social Security Number [SSN], names, dates of birth) to be linked to the NDI. Of the 4.5 million ACS respondents, 33,000 (0.7%) lacked sufficient identifiers and were not sent to the National Vital Statistics System (NVSS) for linkage. An additional 7.6% were sent to NVSS but without SSN. When we analyzed mortality rates based on the linked data, we found that whereas 6.8% of all respondents died during the follow-up period, only 1.2% of respondents without a SSN died during follow-up (**Table S1**). After

---

<sup>10</sup> US Census Bureau. Mortality Disparities in American Communities (MDAC): Project overview. <https://www.census.gov/topics/research/mdac.html>

<sup>11</sup> *Ibid.*

Preliminary. Please do not cite/distribute without permission.

consultation with the Census statistician, we concluded that this most likely reflects under-linkage among persons without a SSN. In the absence of a SSN, the NVSS would have required a very close match on names and date of birth, leading to a lower linkage rate. Based on this analysis, we decided to exclude from the sample respondents without a SSN, because their mortality rates would be underestimated. People without a SSN were somewhat more likely to be poor, less educated, rural, immigrants, and to reside in group quarters. Differences were relatively small, however.

We assessed for evidence of a “healthy respondent” effect in the ACS. We find some suggestive evidence that mortality rates are a bit lower in year 1 for working age adults – consistent with survey non-participation of people in poor health. However, the differences were not very large, were not apparent in other age groups, and we decided not to exclude the first year of follow-up in our analysis (**Table S2**). Another limitation of the linked data is that persons who migrate outside of the U.S. remain in the denominator but would not be observed in U.S. vital statistics. These “immortals” could artificially deflate mortality estimates. We note that external migration is relatively low among AIAN, but it may affect mortality estimates of other groups, particularly those with large immigrant populations.

Overall, mortality rates in the MDAC data matched up with the CDC’s reported mortality rates for the whole U.S. population (**Table S3**). Initial analyses revealed age-adjusted mortality estimates substantially lower than what CDC had reported nationally. (We extracted national data on deaths and population denominators for the period 2008-2015 from CDC WONDER as a comparison.) Excluding people without SSNs made a substantial difference, bringing the mortality rates in the MDAC much closer to the CDC reported national data. **Table S3** shows that MDAC age-specific mortality rates line up closely with the national CDC data at ages 5-34 and ages 65+, with a modest undercount observed at ages 35-64. Mortality rates in the first year of life are severely underestimated. Whereas the national infant mortality rate was 617 per 100K, the MDAC estimate was 111 per 100K. It is likely that the ACS simply misses newborns who die before they leave the hospital. Additionally small numbers of deaths among children mean that data do not meet Census suppression rules for release. Therefore, we focus on the population ages 15 and older.

### *Study Sample*

After excluding people without SSNs, our sample included 4,136,000 ACS respondents, including 58,000 people who self-identified as American Indian or Alaskan Native (AIAN). We defined AIAN as non-Hispanic persons reporting AIAN alone or in combination with other racial categories. **Table 1** shows the characteristics of the study sample. AIAN had similar poverty and education levels as Black and Hispanic respondents but were much more likely to reside in rural areas than those demographic groups. 24% of AIAN resided in a Census-defined “AIAN area” including an Indian Reservation or other designated area.

### **3. Misclassification of AIAN on death certificates**

**Table 1** shows the deaths occurring in the linked MDAC database. We observed 58,000 people who self-identified as AIAN for 423,000 person years of follow-up, during which period 926 self-identified AIAN died. When relying on race/ethnicity as reported in death certificates, however, only 449 of all deaths in the MDAC database had “AIAN” recorded as the decedent’s race. In other words, using death certificate-based race/ethnicity instead of self-reported race/ethnicity would yield death counts (and mortality rates) that were 48.5% of what they should be, implying that true AIAN mortality could be as much as twice as high as existing unadjusted estimates. This number (48.5%) is substantially lower than the 75% estimated by Arias et al. using 2010 Census data (Arias 2021) and we are working to resolve this discrepancy.

**Table 2** shows the racial classification matrix of deaths observed in the MDAC database. Nearly 60% of deaths occurring to self-identified AIAN were misclassified on death certificates, with 51% coded as non-Hispanic White and 7% coded as non-Hispanic Black. Deaths among self-identified AIAN were much more likely to be misclassified than any other group, with just 0.5% of White decedents misclassified, 3% of Black decedents misclassified, 7% of Asian or Pacific Islander decedents misclassified, and 13% of Hispanic decedents misclassified, compared to 60% of AIAN decedents.

#### 4. Impact of misclassification on death rates

**Table 1** shows crude death rates for AIAN when using self-identified vs. death certificate assigned race/ethnicity. When using self-identified race/ethnicity, the crude death rate was 926 per 100K, the second-highest of any racial group. When using death certificate assigned race/ethnicity, the crude death rate was 449, about 50% lower and the second-lowest of any racial group. Of course, difference across racial groups are driven in part by differences in the age distribution. **Table 3** shows results by age group. The table compares death rates for the AIAN population when using self-identified race/ethnicity vs. race/ethnicity from death certificates. The table also reports misclassification ratios by age. Relying on death certificate-assigned race leads to underestimating death rates under 45 years by about one third and over 45 years by over half. To take one example, the estimated death rate for AIAN ages 65-74 using death certificate race/ethnicity was 1180 per 100K. When using self-reported race, the death rate was 2720 per 100K.

One reason for the large observed difference is that the race data on death certificates are provided with “bridged race” categories, in which people who report multiple races are assigned to a single race. Some people who self-reported AIAN and another category may have been bridged into the other category on the death certificate. We therefore conducted additional analyses comparing the MDAC self-report data with CDC Wonder data which use bridged race categories in both the numerator and denominator.

**Figure 1** displays the age-specific mortality rates shown in Table 3 alongside the equivalent estimates from CDC Wonder (2008-2015), calculated using separate numerator and denominator data. The yellow line in Figure 1 shows the ratio of AIAN mortality as estimated in CDC Wonder data vs. our MDAC estimates. The ratio declines with age, indicating that CDC Wonder data are more complete for younger age groups and the impact of misclassification is larger at older age groups.

Preliminary. Please do not cite/distribute without permission.

**Figure 2** shows mortality rate ratios comparing AIAN age-specific mortality rates with age-specific mortality rates for the whole U.S. population regardless of race/ethnicity. The line with solid circles shows CDC Wonder data. The line with hollow circles shows MDAC data, which lie above the CDC Wonder data for all age groups. AIAN death rates are more than twice the national average for persons ages 15-54 and remain above the national average at all age groups. The survival advantage previously reported among older AIAN based on CDC Wonder data may be an artifact of misclassification.

**Figure 3** summarizes the results across age groups in age-standardized mortality rates. Across all racial/ethnic groups, age standardized mortality in the MDAC sample was 1063 deaths per 100,000. The mortality rate among AIAN was 1324 per 100,000 when using race as reported on death certificate and 1463 per 100,000 when using self-reported race. The gap between AIAN age-standardized mortality and total age-standardized mortality increases by 53% after accounting for misclassification.

## 5. Explaining the AIAN mortality disadvantage

A final aim of our analysis was to assess what mediating factors are associated with excess mortality among AIAN relative to other U.S. population groups. **Table 4** shows preliminary results. We fit Cox Proportional Hazards models to assess whether being self-identified Non-Hispanic AIAN is associated with higher (or lower) mortality. The models all control flexibly for age. Columns (1) and (3) compare non-Hispanic AIAN to non-Hispanic Whites. Columns (2) and (4) compare non-Hispanic AIAN to all other groups. Columns (1) and (2) show that being AIAN was associated with about a 40% increased mortality risk across follow up. Columns (3) and (4) show that adjusting for economic, social, and demographic factors (see Table for list of controls included) reduced this association by about half to around 20% increased risk. In other words, about half of the excess mortality risk is statistically explained by factors such as living in a rural area and having lower income / education / employment.




Preliminary. Please do not cite/distribute without permission.

## **6. Conclusion**

The AIAN mortality disadvantage relative to other U.S. racial/ethnic groups is substantially larger than it appears in traditional CDC estimates. Addressing high mortality rates among AIAN – a legacy of past and present discrimination and disadvantage – is an urgent policy priority.


## FIGURES

**Figure 1. Age-specific mortality rates for AIAN: CDC Wonder, MDAC, ratio**




Preliminary. Please do not cite/distribute without permission.

**Figure 2. AIAN mortality relative to total U.S. mortality by age: CDC Wonder vs. MDAC**



Preliminary. Please do not cite/distribute without permission.

**Figure 3. Age-standardized mortality rates (ages 15+)**



Preliminary. Please do not cite/distribute without permission.

## TABLES AND FIGURES

**Table 1. Description of the Study Sample, by Self-Reported Race and Ethnicity**

Parameter	NH AIAN	NH White	NH Black	NH API	Hispanic	US Total
<i>Sample Size (unweighted counts)</i>						
People	58,000	3,072,000	396,000	187,000	424,000	4,136,000
Person-years (based on ACS race)	423,000	22,200,000	2,880,000	1,380,000	3,140,000	30,000,000
Deaths (based on ACS race)	3900	253,000	26,500	7100	13,500	304,000
Deaths/100,000 PY (ACS race)	926	1141	916	519	425	1013
Deaths (based on DC race)	1900	257,000	26,500	6800	12,500	304,000
Deaths/100,000 (DC race)	449	1156	920	494	395	1013
Ratio: DC deaths to ACS deaths	0.485	1.013	1.005	0.953	0.929	1.00
<i>Respondent Characteristics</i>						
% female	51.2%	50.9%	52.6%	51.4%	49.6%	51.0%
Mean age (years)	33.6	40.2	33.2	34.8	29.3	37.5
% <FPL	20.4%	8.6%	22.3%	8.9%	18.5%	11.8%
% rural	36.3%	29.2%	12.3%	8.2%	9.8%	23.6%
% college	10.8%	22.9%	11.3%	34.2%	8.4%	19.9%
% in "AIAN area"	24.3%	1.6%	1.8%	0.7%	0.8%	1.8%

Note: Sample is the 2008 ACS, restricted to those participants with a SSN available for linkage to the National Death Index. Deaths are those that were linked to the 2008 ACS sample. Respondent characteristics are as assessed in the 2008 ACS and are weighted to be representative of the U.S. resident population. "AIAN area" is a Census designation Census blocks or tracts that are contained within American Indian or Alaska Native Legal or Statistical Areas, including American Indian Reservations, Alaska Native Village Statistical Areas, and Oklahoma Tribal Statistical Areas (Census 2021, Understanding and Using ACS Data: What Users of Data for American Indians and Alaska Natives Need to Know).

Preliminary. Please do not cite/distribute without permission.

**Table 2. Classification of race and ethnicity in death certificates, by self-reported race and ethnicity**

ACS Race/ethnicity	% of deaths assigned to each racial / ethnic group on death certificate					Total # of deaths
	NH AIAN	NH White	NH Black	NH API	Hispanic	Total
ACS NH AIAN	40.5%	51.3%	6.7%	0.3%	1.1%	3,900
ACS NH White	0.1%	99.5%	0.2%	0.0%	0.3%	253,000
ACS NH Black	0.1%	2.1%	97.3%	0.1%	0.3%	26,500
ACS NH API	0.1%	5.9%	0.6%	92.7%	0.7%	7,100
ACS Hispanic	0.3%	11.6%	1.0%	0.3%	86.8%	13,500

Note: Table shows data on 304,000 deaths observed in the MDAC study sample. This includes all linked deaths for all 2008 ACS respondents with sufficient information to be sent to NDI for linkage (failedit==0) and who had a social security number.

Preliminary. Please do not cite/distribute without permission.

**Table 3. Age-specific mortality rates for non-Hispanic AIAN (alone or in combination) during 8-year follow-up**

Age	(A) Person-Years	Race/ethnicity from death certificates		Race/ethnicity from ACS self-report		Bias due to misclassification of race/ethnicity in death certificates	
		(B) Deaths	(C) Rate	(D) Deaths	(E) Rate	Ratio (C / E)	Difference (C – E)
Under 1y	400	- S -	- S -	- S -	- S -	- S -	- S -
1-4 years	10000	- S -	- S -	- S -	- S -	- S -	- S -
5-14	62500	- S -	- S -	- S -	- S -	- S -	- S -
15-24	72500	70	89.5	90	125	0.714	(-35.8)
25-34	53500	90	161	150	236	0.683	(-74.8)
35-44	52500	100	208	200	367	0.568	(-159)
45-54	63500	200	329	450	675	0.488	(-345)
55-64	56500	300	548	650	1150	0.478	(-599)
65-74	32500	400	1180	900	2720	0.433	(-1540)
75-84	14500	400	2910	850	5970	0.487	(-3060)
85+	4700	300	6580	650	14400	0.458	(-7780)

Note: Mortality rates were based on the 2008 ACS sample that had social security numbers available for linkage to the National Death Index. (A) Follow-up time (person-years) was defined as being from the date of the 2008 ACS interview to Dec 31, 2015, or to the date of death as identified in the NDI match. (B) Deaths (ACS race) are those that were linked to the 2008 ACS records for non-Hispanic AIAN. (C) Deaths (DC race) are the count of deaths that occurred to persons who were identified as non-Hispanic AIAN in the death certificate, from the sub-sample of linked deaths. (D) The death rate *estimates* (per 100,000 person-years) are based on unweighted deaths and person-years. (E) “- S -” indicates the count or estimated death rate is suppressed due to fewer than 15 deaths supporting the estimate.

Preliminary. Please do not cite/distribute without permission.

**Table 4. Cox proportional hazards models: Estimates hazard ratios for risk of death during 8-year follow-up**

<b>Hazard ratio (95% CI)</b>	<b>(1)</b>	<b>(2)</b>	<b>(3)</b>	<b>(4)</b>
<i>Race/ethnicity</i>				
NH AIAN	1.425	1.422	1.260	1.223
NH White	Ref	Ref	Ref	Ref
NH Black	Ref	1.237	Ref	1.068
NH API	Ref	0.795	Ref	0.779
Hispanic	Ref	0.792	Ref	0.664
<i>Socio-demographics</i>				
Income				
< Poverty rate			1.148	1.163
1.00-1.99 * Pov.Rate			1.043	1.054
2.00-2.99 * Pov.Rate			0.975	0.983
3.00-4.99 * Pov.Rate			0.915	0.920
> 5 * Pov.Rate			Ref	Ref
Education				
Dropout			1.109	1.158
H.S. Diploma			Ref	Ref
Some College			0.863	0.860
College Graduate			0.670	0.670
Employment Status				
Employed			0.734	0.734



Preliminary. Please do not cite/distribute without permission.

Unemployed			1.305	1.313
Not In Labor Force			1.632	1.644
Other (children, Armed Forces)			Ref	Ref
Rural			0.965	0.935
In AIAN area			1.128	1.110
Immigrant (in last 5 years)	0.643	0.726	0.573	0.670
<i>Years since ACS interview</i>				
0 to <1	1.043	1.040	0.968	0.961
1 to <2	1.037	1.035	0.976	0.970
2 to <3	1.041	1.040	0.993	0.989
3 to <4	1.010	1.009	0.976	0.974
4 or more	Ref	Ref	Ref	Ref

---

Note: All coefficients are significant at  $\alpha=0.05$ . All standard errors for coefficients are less than 0.008. Table presents Cox-proportional hazards models using age as the time scale. In other words, age is adjusted for non-parametrically in all models. Sample is the 2008 ACS sample that had social security numbers available for linkage to the National Death Index.

Preliminary. Please do not cite/distribute without permission.

**Table S1. Supporting Data for Defining the Study Sample, 2008-2015 follow-up**

<b>Parameter</b>	<b>Respondents (N)</b>	<b>Deaths (N)</b>	<b>% died</b>
<i>Non-Hispanic AIAN</i>			
2008 ACS respondents	66,000	4000	6.1%
Not sent to NDI for linkage (failedit ==1)	500	N/A	N/A
Sent to NDI for linkage (failedit ==0), NO Social Security Number	7600	100	1.4%
Sent to NDI for linkage (failedit ==0) with Social Security Number	58,000	3900	6.8%
Immigrated to the U.S. in the prior 5 years	150	- S -	0.0%
<i>All racial and ethnic groups</i>			
2008 ACS respondents	4,510,000	308,000	6.8%
Not sent to NDI for linkage (failedit ==1)	33,000	N/A	N/A
Sent to NDI for linkage (failedit ==0), NO Social Security Number	344,000	4100	1.2%
Sent to NDI for linkage (failedit ==0) with Social Security Number	4,140,000	304,000	7.4%
Immigrated to the U.S. in the prior 5 years	66,500	750	1.1%

Note: “- S -” indicates that the count was suppressed for being below 15.

Preliminary. Please do not cite/distribute without permission.

**Table S2. Age-specific mortality, by year of follow-up (all racial and ethnic groups, 2008-2015)**

Parameter	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7
<b>Deaths per 100K person years</b>							
Under 1 year	111	.	.	.	.	.	.
1-4 years	24	20	.	.	.	.	.
5-14	14	14	13	15	13	13	.
15-24	66	67	62	66	72	62	66
25-34	93	99	102	101	105	97	114
35-44	151	167	167	155	167	158	176
45-54	358	372	377	362	355	361	363
55-64	777	785	790	781	802	776	765
65-74	1857	1846	1853	1785	1778	1756	1738
75-84	4790	4780	4864	4697	4664	4659	4682
85+	14110	13600	13880	13370	13960	13340	13940

Note: *Estimates* of mortality rates are based on person-time contributed to that age group during the 8-year follow-up, depending upon the respondent's age at the beginning of each person-year (so respondents can age into older groups). Data are restricted to ACS respondents with a social security number, submitted to the NDI for linkage.

Preliminary. Please do not cite/distribute without permission.

**Table S3. Age-specific mortality 2008-2015 in MDAC sample vs. CDC WONDER**

<b>Deaths per 100K</b>	<b>MDAC</b>	<b>MDAC</b>	<b>CDC WONDER</b>
<b>Age group</b>	<b>(full sample)</b>	<b>(after exclusions)</b>	<b>(pooled, 2008-2015)</b>
Under 1 year	99	111	617
1-4 years	18	19	26
5-14	13	14	13
15-24	61	66	68
25-34	94	102	107
35-44	152	163	175
45-54	343	363	409
55-64	742	781	861
65-74	1702	1790	1839
75-84	4469	4714	4726
85+	12,540	13,710	13,750

Note: MDAC full sample is the 2008 ACS. MDAC after exclusions is the 2008 ACS after excluding: respondents with insufficient data to be sent to the NDI for linkage; respondents without a social security number.